

Exploring different definitions of methane phenotypes in Dutch Holstein Cattle

C. I. V. Manzanilla-Pech,^{1*} A. E. van Breukelen,¹ R. F. Veerkamp,¹ Y. de Haas,¹ M. van Pelt,² and B. Gredler-Grandl¹

Abstract One of the most promising strategies to permanently reduce methane emissions in dairy cattle is through genomic selection, where the primary goal is to identify and selectively breed low-emitting ruminants. An important step is to define which trait definition to use. Several methane phenotypes have been proposed in recent years. When measuring methane using breath analyzers (sniffers), 2 commonly used phenotypes are methane concentration (CH₄c; ppm) and methane production (CH₄p; g/d). However, for both phenotypes different definitions exist. For example, CH₄c, the average per visit is commonly used, but other phenotypes involving the identification of “eructation” peaks have been suggested. For CH₄p several formulas are available to calculate it, using CH₄c or the ratio between CH₄c and CO₂ concentrations as input, together with other production traits. Additionally, methane intensity, another phenotype of interest, is the amount of methane produced per milk unit (kg). Therefore, the aims of this study were to: 1) Estimate genetic parameters for 11 distinct phenotypes, including 3 CH₄c definitions, carbon dioxide concentration per visit (CO₂c), the ratio between CH₄c and CO₂c, 5 phenotypes for CH₄p based on different formulas (Madsen, Chagunda, 2 of Kjeldsen, and IPCC-Tier2 as benchmark), and methane intensity (MeI). 2) Estimate genetic correlations between these methane phenotypes and milk yield (MY) and body weight (BW). A total of 149,726 sniffer (CH₄c and CO₂c) records were available from 7,600 Dutch Holstein cows measured between 2019 and 2024 on 68 farms. Data were analyzed with an animal repeatability model with fixed effects, including herd-year-season, week of lactation and lactation number with age of cow at calving nested. Estimated heritability values ranged between 0.16 and 0.30 for CH₄c phenotypes, with the number of eructation peaks having the highest heritability. For CH₄p, heritabilities ranged between 0.03 to 0.27, with Tier2 having the highest value. Heritabilities for ratio and CO₂c were 0.08 and 0.13, respectively. Genetic correlations between CH₄c phenotypes were moderate to highly positive, ranging between 0.49 and 0.85. Likewise, highly positive genetic correlations (between 0.89 and 1) were estimated for CH₄p phenotypes, except for Tier2, which presented correlations between 0.04 to 0.37 with the other CH₄p phenotypes. Both mean CH₄c (avg) and sum of max peaks for CH₄c (speaks) had moderate to high positive correlations (0.36–0.95) with the majority of the CH₄p phenotypes (except Tier2). Methane intensity had moderate to high positive genetic correlations (0.38 to 0.80) with the majority of the phenotypes except for Kjeldsen2 (0.03) and Tier2 (−0.90). Milk yield had positive genetic correlations with all the methane phenotypes (0.04 to 0.94) except for MeI (−0.66) and Kjeldsen2 (−0.69). Body weight had close to zero genetic correlations with CH₄c phenotypes (−0.09 to 0.07), and moderate positive to moderate negative correlation (−0.72 to 0.57) for CH₄p phenotypes. Given their strong correlations with the other methane phenotypes, close to zero correlation with body weight, and no induced dependencies with BW and MY, as seen with the CH₄p phenotypes, avg and speaks appear to suitable proxies for methane emissions when using sniffers.

Methane emissions in dairy cattle have been investigated intensively over the past decade due to the 2030–2050 climate targets, where the EU aims for a 55% reduction up to neutrality in greenhouse gas emissions (EEA, 2023). One of the mitigation options is through genetic/genomic selection, where the primary goal is to identify and selectively breed low-emitting animals. As a result, several countries are currently monitoring enteric methane from ruminants. A widely employed phenotyping method involves breath sampling during milking by a breath analyzer device commonly referred to as “sniffer.” This device samples breath at regular intervals (often between 1 and 15 s) during milking, reporting methane concentration (CH₄c) in parts per million (ppm). While CH₄c has been proposed as an indicator for gross methane emissions (referred to as methane production (CH₄p); CH₄ g/d) due to its high correlation (0.76; van Breukelen et al., 2022), there is a lack of consensus on which phenotype to use for estimating breed-

ing values. Several phenotypes have been proposed over the past years; one largely used is the average CH₄c per visit or per minute averaged daily or weekly, but other CH₄c phenotypes involving the identification of “eructation” peaks have also been suggested (Garnsworthy et al., 2012; Rey et al., 2019; Reintke et al., 2020). Both studies that investigated peaks used CH₄c measured by mobile laser methane detectors (LMDs). Rey et al. (2019) defined the peaks as the number of peaks or eructation events presented in 5 min. Reintke et al. (2020) proposed the sum of CH₄c per minute during eructation peaks, maximum CH₄c during eructation peaks, and the number of eructation events per minute as possible CH₄c phenotypes in ewe breeding. The rationale behind the use of eructation peaks as a phenotype is to try to disentangle CH₄c that comes from a respiration event from the one that comes from an eructation event (i.e., the peaks). Previous studies by nutritionists (Blaxter and Joyce, 1963 and Murray et al., 1976) showed that

¹Animal Breeding and Genomics, Wageningen Livestock Research, Wageningen University & Research, 6708 PB Wageningen, the Netherlands, ²Cooperation CRV, Animal Evaluation Unit, P.O. Box 454, 6800AL Arnhem, Netherlands. *Corresponding author: coralia.manzanillapech@wur.nl. © , The Authors. Published by Elsevier Inc. on behalf of the American Dairy Science Association®. This is an open access article under the CC BY license (<https://creativecommons.org/licenses/by/4.0/>). Received July 27, 2025. Accepted September 29, 2025.

The list of standard abbreviations for JDSC is available at [adsa.org/jdsc-abbreviations-25](https://www.adsa.org/jdsc-abbreviations-25). Nonstandard abbreviations are available in the Notes.

approximately 17% of the methane exhaled originated from the lungs, while the remaining 83% was produced by eructation. Additionally, eructation peak detection could adjust for barn ambient CH₄c, but also for build-up of CH₄c in the AMS feed bin during milking (Bell, et al., 2014). While there is no published literature on the heritability of eructation peaks in cows, several studies have reported the detection of methane peaks using sniffers (Rey et al., 2019; Hardan et al., 2022). Additionally, sniffers traditionally rely on formulas to “convert” CH₄c in ppm to CH₄p in g/d. Currently, there are several formulas available to calculate CH₄p using CH₄c and body weight (BW; Chagunda et al., 2009) or predicting carbon dioxide (CO₂ g/d; via heat production using BW, energy-corrected milk, and days in pregnancy) and multiplying it by the ratio between CH₄c and CO₂c to predict CH₄p (Pedersen et al., 2008; Madsen et al., 2010). Despite some of these formulas being extensively used, they were originally developed to predict CH₄p in barns, not individual animal CH₄p. Recently, Kjeldsen et al. (2024) developed 3 formulas to predict CO₂ g/d using an international data set of more than 2 thousand cows with recorded CH₄p using either respiration chambers or GreenFeed. Finally, the IPCC (2019) has proposed several approaches (Tier1 to 3) to calculate methane emission factors per species, where the Tier2 formula is extensively used for dairy cattle, as it uses an energy balance approach. Therefore, the aims of this study were to: 1) Estimate genetic parameters for 11 distinct phenotypes, including 3 CH₄c definitions, CO₂ concentration (CO₂c) per visit, the ratio between average CH₄c and CO₂c, 5 phenotypes for CH₄p based on different formulas (Madsen, Chagunda, Kjeldsen, and Tier2), and methane intensity (MeI); and 2) Estimate genetic correlations between these methane phenotypes and milk yield (MY, kg) and BW.

The original data included a total of 149,726 sniffer CH₄c and CO₂c records from roughly 7,600 primi- and multiparous Dutch Holstein cows. Data has been previously described in van Breukelen et al. (2023; 2024). Methane concentration was measured using sniffers located in automated milking systems (AMS) (WD-WUR v2.0, Carltech BV), sampling CH₄c and CO₂c every 5 s. Based on the sniffer CH₄c and CO₂c (ppm) measurements, 11 phenotypes were derived. We divided the phenotypes into groups: CH₄c in ppm, CH₄p in g/d and methane intensity (MeI in g CH₄/kg MY). Methane concentration phenotypes included 1) average CH₄c per visit (**avg**), 2) number of eructation peaks per minute (**npeaks**), 3) sum of maximum eructation peaks defined as the sum of the average of the 2 top values within each peak (for all peaks within visit; **speaks**). Methane production phenotypes were calculated with 5 formulas available in the literature: 4) **Madsen** (Madsen et al., 2010) based on the prediction of CO₂ using ECM and BW to posteriorly multiply it by ratio; 5) **Chagunda** (Chagunda et al., 2009) based on the tidal volume that uses BW as a predictor and the average CH₄c; 6) and 7) **Kjeldsen2** and **Kjeldsen3** (Kjeldsen et al., 2024) based on the prediction of CO₂ using ECM and BW or only ECM, and multiplied by the ratio between CH₄c and CO₂c; 8) **Tier2** (IPCC, 2019) based on the prediction of CH₄p using MY and BW. Tier 2 was used as a benchmark scenario incorporating information on MY, body weight BW, and diet, without relying on sniffer-based measurements, and was not treated as a selection phenotype. Additionally, we included as phenotypes 9) the **ratio** of CH₄c to CO₂c, 10) CO₂ concentration (**CO₂c**), and 11) methane intensity (**MeI**) defined as CH₄p divided by MY (g CH₄p/kg MY), where CH₄p was calculated using the Kjeldsen3 formula (as it was

the phenotype with the largest number of records and animals). Each of the 11 phenotypes were calculated at visit level. Weekly CH₄c and CH₄p phenotypes were calculated averaging visit phenotypes (with a minimum number of 5 visits) per calendar week. All CH₄c phenotypes had a background level subtracted, where the background was calculated as the 5 lowest measurements during the entire visit, with the purpose to account for the CH₄c level in the barn. Furthermore, visit CH₄c and CH₄p was defined as the concentration present in a window of 240 s (between 60 and 300 s after the entrance time of the cow in the AMS). The first 60 s were removed to account for the gas traveling through the tube until the gas sensor. Visits were cut at 300 s and visits with less than 300s were discarded to avoid erratic data due to head movement as shown by Løvendahl, et al. (2024), as pellets are dropped in the AMS and consumed (mainly) in the first minutes of the milking. Eructation peaks were defined using a self-written function in R to detect peaks. This approach detects local maxima by computing the first and second differences of the signal (0.0005). The function calculates the change between consecutive values. A peak is defined as a point where the signal transitions from increasing to decreasing — i.e., where the first derivative is positive followed by a negative slope. Once eructation peaks were defined, 2 phenotypes were calculated, a) number of peaks per minute, that was the total number of peaks per visit divided by visit length and b) the 2 maximum values per peak were detected, averaged and then summed across all the peaks per visit. The majority of animals had MY and milk content information, however, only ~10% of the animals had BW. For this reason, phenotypes that used BW or metabolic BW (MBW) had a smaller number of records and animals. The number of records and animals per phenotype after editing is presented in Table 1.

The CH₄p phenotypes were calculated using the following formulas:

$$\text{Madsen: CO}_2 \text{ (g/d)} = 180 \times 24 \times (5.6 \text{ MBW} + 22 \text{ ECM} + 1.6 \times 10^{-5} \times \text{num days in gestation})$$

$$\text{Chagunda: CH}_4\text{p (g/d)} = \text{mean CH}_4\text{c}/10^6 \mu\text{L} \times \text{tidal volume(L/ breath)} \times 30 \text{ breaths/min} \times 1,440 \text{ min/d} \times 16.04 \text{ (g/mol)}/22.4 \text{ (L/mol)}$$

where TV(L) = 7.5MI / kgofLW × LW(kg)/1,000(mL/L)

Kjeldsen2

$$\begin{aligned} \text{CO}_2 \text{ (g/d)} = & -6134 + (213 \times \text{ECM}) + (126 \times \text{MBW}) + (52.5 \times \\ & \text{Milk CF}) + (-5.13 \times \text{DIM}) + 2117 + (-0.122 \times \text{DIM} \times \text{Diet CF}) \\ & + (0.386 \times \text{ECM} \times \text{DIM}) + (-1.18 \times \text{ECM} \times \text{MBW}) + (-0.614 \times \\ & \text{Milk CF} \times \text{MBW}) + (-5.96 \times \text{MBW}) + (-1.18 \times \text{DIM}) + (-0.614 \\ & \times \text{MBW}) \end{aligned}$$

where Milk CF is crude fat in milk, Diet CF is crude fat in the diet.

Kjeldsen3

$$\begin{aligned} \text{CO}_2 \text{ (g/d)} = & 8781 + (80.3 \times \text{ECM}) + (-4.66 \times \text{DIM}) + -49.0 \times \\ & \text{parity coefficient} + \text{breed-parity coefficient} + (-0.149 \times \text{DIM} \times \\ & \text{Diet CF}) + (0.338 \times \text{ECM} \times \text{DIM}) + (206 \times \text{DIM}) + (\text{Milk CF} - \end{aligned}$$

parity coefficient \times Milk CF)

where parity coefficients (Kjeldsen et al., 2024) are 0 for first, 511 for second and 1587 for third and higher parities, breed-parity coefficients are for Holstein 0 for first, 775 for second and 803 for third and higher, and MilkCF-parity coefficients are -4.18 for first, -10.5 for second and -28.8 for third and higher.

Tier2 (Equation 10.21)

$$\text{CH}_4\text{p (g/d)} = (\text{GE} * 0.065 / 55.65) * 1000$$

$$\text{GE} = (\text{NEm} + \text{NEa} + \text{NE}_1 + \text{NEp}) / \text{REM} / \text{DE} / 100, \text{REM} = 1.123 - (0.004092) * \text{DE} + 0.00001126 * (\text{DE} - 25.4 / \text{DE}) \text{ and CF} * \text{MBW}$$

where gross energy (GE) was calculated with the (Equation 10.16) ignoring the net energy (NE) for grow (as they are grown animals and the NE for wool. The $\text{NE}_1 = \text{Milk} * (1.47 + 0.4 * \text{Kg Fat})$ and NEm is net energy for maintenance, NEa is net energy for activity, NE_1 is net energy for lactation, REM is ratio of net energy available in a diet for maintenance to digestible energy, DE is digestible energy as percentage of gross energy and CF is crude fat.

Madsen and Kjeldsen's formulas predicted CO_2 g/d, and this prediction was multiplied by the ratio of CH_4c to CO_2c to calculate CH_4p . Variance components were estimated with a repeatability animal model in ASReml v. 4.2.1. The model was as follows:

$$\mathbf{y} = \mathbf{Xb} + \mathbf{Z_a a} + \mathbf{Z_p pe} + \mathbf{e},$$

where \mathbf{y} is the vector of phenotypes; \mathbf{b} is the vector of fixed effects: herd, year-season interaction ($n = 145$), week of lactation (1–60), age of cow at calving in months (21–122) nested within parity (1, 2, ≥ 3); \mathbf{a} is the vector of direct additive genetic effects; \mathbf{pe} the vector of random permanent environment effects; and \mathbf{e} is the vector of residual effects. The matrices \mathbf{X} , $\mathbf{Z_a}$ and $\mathbf{Z_p}$ are the incidence matrix relating observations with the fixed effects, random genetic effects, and random permanent environment effects, respectively. Distributions of the random effects are (1) $\text{var}(\mathbf{a}) = \mathbf{A}\sigma_a^2$ where \mathbf{A} is the pedigree relationship matrix and σ_a^2 is the animal additive genetic variance, and (2) $\text{var}(\mathbf{pe}) = \mathbf{I}\sigma_{pe}^2$, where \mathbf{I} is an identity matrix of

an order equal to the number of observations and σ_{pe}^2 is the permanent environmental variance, and (3) $\text{var}(\mathbf{e}) = \mathbf{I}\sigma_e^2$, where \mathbf{I} is an identity matrix of an order equal to the number of observations and σ_e^2 is the residual variance. Pedigree included 64,334 animals with on average 14 generations. Bivariate analyses were conducted to estimate the genetic correlations among the different phenotypes. Due to the imbalance in record numbers of animals and records between traits, estimates of some pairwise bivariate analyses may be less precise and have reported large standard errors. Therefore, genetic parameters involving these traits (Madsen and Kjeldsen 2) should be interpreted with caution.

Descriptive statistics for the different phenotypes are summarized in Table 1. The average CH_4c phenotypes were 401.8 ppm for conc, 1.5 for npeaks, and 954.1 ppm for speaks. The average CH_4p (g/d) varied depending on the formula, ranging from 189.0 ± 101.8 (Chagunda) to 355.1 ± 68.1 (Tier2). The average CH_4p (g/d) was 296.1 ± 132.0 for Madsen, 309.3 ± 135.1 for Kjeldsen2, and 318.5 ± 128.8 for Kjeldsen3, indicating that these formulas predicted similar values for CH_4p , which was expected as their formulas are the most similar. The average MeI was 9.9 g CH_4p per kg of MY.

Estimated variance components are presented in Table 2. Heritabilities for CH_4c phenotypes were 0.16 for conc, 0.29 for speaks, and 0.30 for npeaks. Heritabilities for average CH_4c have been widely reported in the literature, ranging between 0.11 and 0.18 (Lopez-Paredes et al., 2020; Manzanilla-Pech et al., 2020; van Breukelen et al., 2023). Unlike these studies, Reintke et al. (2020) was the only study reporting heritability for the number of eructation peaks, which was 0.02 with a standard error (SE) of 0.05. This study was conducted in ewes, and the lower heritability may be due to the method (laser methane detector) rather than the phenotype. Heritabilities for CH_4p ranged between 0.03 (Madsen) and 0.27 (Tier2). The heritabilities for the ratio and MeI were 0.08, and was 0.13 for CO_2c . Larger heritabilities (0.11–0.47) have been reported for CH_4p using the Madsen formula by Manzanilla et al. (2022a, 2023) in a Danish Holstein population and by Sypniewski et al. (2021) in a Polish Holstein population. Genetic correlations between CH_4c phenotypes (Figure 1) ranged from moderate between avg and npeaks (0.49, SE = 0.06) to highly positive between avg and speaks (0.85, SE = 0.02). As expected, the number of eructation peaks is moderately correlated with the 2 other CH_4c phenotypes. However, since it does not reflect the intensity of the eructation,

Table 1. Descriptive statistics for the CH_4c per visit (avg), sum of maximum peaks per minute (speaks), number of peaks per minute (npeaks), CH_4p calculated by Madsen, Chagunda, Kjeldsen2, Kjeldsen3 and Tier2, ratio between CH_4c and CO_2c , carbon dioxide per visit (CO_2c), methane intensity (MeI), milk yield (MY) and body weight (BW)

Type	Trait	#cows	#records	mean	SD	CV (%)
CH_4c (ppm)	avg	7,110	114,574	401.8	179.1	44
	npeaks	7,129	114,744	1.5	0.4	26
	speaks	7,018	114,244	954.1	458.9	48
CH_4p (g/d)	Madsen	806	10,668	296.1	132.0	44
	Chagunda	868	16,822	189.0	101.8	54
	Kjeldsen2	798	10,676	309.3	135.1	44
	Kjeldsen3	6,608	73,799	318.5	128.8	40
	Tier2	855	14,635	355.1	68.1	19
	ratio	6,945	99,931	0.07	0.02	28
	CO_2c	7,529	141,591	5,376.0	1,846.0	34
Intensity (g CH_4p / kg MY)	MeI	6,608	73,799	9.9	4.8	48
	MY (kg/d)	6,774	84,287	34.4	9.6	28
Production traits	BW (kg)	868	16,868	697.6	82.7	12

Table 2. Estimated genetic variance (σ_a^2), phenotypic variance (σ_p^2), heritability (h^2) and repeatability (rep) for the average CH₄c per visit (avg), sum of maximum peaks per minute (speaks), number of peaks per minute (npeaks), CH₄p calculated by Madsen, Chagunda, Kjeldsen2, Kjeldsen3 and Tier2; carbon dioxide concentration per visit (CO₂c), milk yield (MY), methane intensity (MeI) and body weight (BW)

Trait	σ_a^2	σ_{pe}^2	σ_p^2	h^2	rep
avg	3540.4	6049.40	21979.0	0.16 (0.02)	0.43 (0.01)
speaks	49496.8	37027.6	171040.0	0.29 (0.02)	0.51 (0.02)
npeaks	567.7	425.558	1889.2	0.30 (0.02)	0.52 (0.02)
Madsen	245.1	2585.2	8369.8	0.03 (0.03)	0.34 (0.03)
Chagunda	1670.0	2548.9	7914.0	0.21 (0.06)	0.53 (0.05)
Kjeldsen2	224.1	2377.39	8062.0	0.03 (0.03)	0.32 (0.03)
Kjeldsen3	707.6	1504.6	8465.7	0.08 (0.01)	0.26 (0.01)
Tier2	704.8	1066.2	2608.7	0.27 (0.07)	0.68 (0.06)
ratio	0.28	0.46	3.53	0.08 (0.01)	0.21 (0.01)
CO ₂ c	2649.1	6448.6	20725.0	0.13 (0.02)	0.44 (0.01)
MY	12.2	22.1	54.6	0.22 (0.02)	0.62 (0.02)
MeI	1.1	3.1	13.5	0.08 (0.01)	0.31 (0.01)
BW	2109.2	1313.2	3760.1	0.56 (0.08)	0.91 (0.08)

only the number, it is not a good candidate for a breeding goal trait. Conversely, speaks is highly correlated with conc, meaning that an animal with a high CH₄c during the entire period also has a high value during the eructation peaks. Genetic correlations between CH₄p phenotypes were highly positive (0.89 to 1.0), except for Tier2, indicating that despite numerical differences in the mean values, they rank the animals similarly (high and low emitters). Tier2 presented low to moderate (0.04 to 0.37) genetic correlations with the other CH₄p phenotypes. Tier2 is used more as a control, as it does not involve any input from the sniffers but relies entirely on MY and BW. It has been developed to predict CH₄p from a group of animals rather than individually (IPCC, 2006; 2019). However, it is important to mention that due to the low number of animals for some of the formulas that include BW in their calculation, some of the SEs were high (0.06 to 0.30). Currently, there is no published information on the genetic correlations between the methane phenotypes presented in this study for either CH₄c or CH₄p, except for avg and CH₄p with Madsen formula. Genetic correlations between avg and CH₄p were high (0.77–0.93), except for Tier2 (–0.08). Genetic correlations between speaks and CH₄p were positive, ranging from 0.36 (Kjeldsen2) to 0.95 (Chagunda), and negative for Tier2 (–0.19). Similarly, npeaks correlations were positive, ranging from 0.52 (Kjeldsen3) to 0.99 (Kjeldsen2), and negative for Tier2 (–0.39). Genetic correlations between ratio and CH₄c ranged from low negative (–0.18 to –0.08; speaks and conc) to moderate positive (0.59; npeaks). In contrast, the majority of genetic correlations between CH₄p phenotypes and ratio were positive, ranging from 0.35 (Madsen) to 0.98 (Kjeldsen3), except for Tier2 (–0.25). No genetic correlation was estimated between CO₂c and npeaks; however, moderate positive genetic correlations from 0.49 (CO₂c–speaks) to high correlations of 0.84 (CO₂c and avg) were estimated. Genetic correlations between CO₂c and CH₄p phenotypes were mostly moderate positive (0.18–0.58), except for Kjeldsen2 (–0.73) with large SE (0.17).

Methane intensity presented the largest correlations with avg (0.77) and the smallest with npeaks (0.29), whereas for CH₄p phenotypes, the largest positive correlation was with Madsen (0.57) and the smallest with Kjeldsen2 (0.03). However, as CH₄p itself is derived from ECM and/or BW, mathematical dependencies may artificially affect genetic correlations with MeI. Further, the

genetic correlation between MeI and Tier2 was the largest negative (–0.90), which may also be affected by the shared use of MY in the calculation of both traits. Genetic correlations between MY and avg (0.30) and between MY and speaks (0.46) were moderately positive, whereas with npeaks it was close to zero. This means that CH₄c phenotypes such as avg and speaks are not entirely independent of milk. Most of the genetic correlations between MY and CH₄p were positive (0.32–0.94), except for Kjeldsen2 (–0.69) with large SE (0.97).

Finally, genetic correlations between BW and all CH₄c phenotypes were close to zero, meaning that body weight is independent of CH₄c, as expected. For CH₄p, the genetic correlations with BW were negative for Madsen (–0.30) and Kjeldsen2 (–0.72) and Kjeldsen3 (–0.23), which can be explained by the indirect use of BW (in combination with ECM) in the prediction of CO₂ for Madsen and Kjeldsen2. Positive genetic correlations between BW and Chagunda (0.36) and Tier2 (0.57) could be explained by the use of BW in the calculation of CH₄p. Unlike the previously reported positive genetic correlation (0.65) between MeI and BW by Manzanilla-Pech et al. (2021) in an international data set, in this study, this correlation was moderately negative (–0.45). One disadvantage of CH₄p phenotypes using the formulas and MeI is the artificially created dependency on other traits that are used for their calculation, such as ECM and BW.

All phenotypes exhibited varying degrees of heritability. Average CH₄c showed consistently high positive genetic correlations with all other methane phenotypes except for ratio and Tier 2 (benchmark). The CH₄p phenotypes calculated using the formulas also presented strong positive correlations among themselves and with other methane traits. Given these results, avg and speaks appear to be good proxies for methane emissions, especially as they showed minimal correlation with body weight and no induced dependencies as the CH₄p based on formulas. Overall, selecting for low-emitting animals is possible, regardless of the phenotype selected. The aim of this study is not to propose selection indicators, but rather to provide a fair comparison of different methane phenotypes that could potentially be used in future selection strategies for lower methane emissions when recording with sniffers. In this context, methane concentration is considered the reference trait for methane emissions.

References

- Bell, M. J., N. Saunders, R. H. Wilcox, E. M. Homer, J. R. Goodman, J. Craigon, and P. C. Garnsworthy. 2014. Methane emissions among individual dairy cows during milking quantified by eructation peaks or ratio with carbon dioxide. *J. Dairy Sci.* 97:6536–6546. <https://doi.org/10.3168/jds.2013-7889>.
- Chagunda, M. G. G., D. Ross, and D. J. Roberts. 2009. On the use of a laser methane detector in dairy cows. *Comput. Electron. Agric.* 68:157–160. <https://doi.org/10.1016/j.compag.2009.05.008>.
- EAA (European Environment Agency). 2023. Methane emissions in the EU: the key to immediate action on climate change. <https://www.eea.europa.eu/publications/methane-emissions-in-the-eu/methane-emissions-in-the-eu/download.pdf.static>
- Garnsworthy, P. C., J. Craigon, J. H. Hernandez-Medrano, and N. Saunders. 2012. On-farm methane measurements during milking correlate with total methane production by individual dairy cows. *J. Dairy Sci.* 95:3166–3180. <https://doi.org/10.3168/jds.2011-4605>.
- Hardan, A., P. C. Garnsworthy, and M. J. Bell. 2022. Detection of Methane Eructation Peaks in Dairy Cows at a Robotic Milking Station Using Signal Processing. *Animals (Basel)* 12:26. <https://doi.org/10.3390/ani12010026>.
- IPCC. 2019 Refinement to the 2006 IPCC Guidelines for National Greenhouse Gas Inventories. Chapter 10: Emissions from livestock and manure management. IPCC Guidelines for National Greenhouse Gas Inventories Volume 4: Agriculture, forestry and other land use, Intergovernmental Panel on Climate Change (2019). https://www.ipcc-nggip.iges.or.jp/public/2019rf/pdf/4_Volume4/19R_V4_Ch10_Livestock.pdf
- Kjeldsen, M. H., M. Johansen, M. R. Weisbjerg, A. L. F. Hellwing, A. Bannink, S. Colombini, L. Crompton, J. Dijkstra, M. Eugène, A. Guinguina, A. N. Hristov, P. Huhtanen, A. Jonker, M. Kreuzer, B. Kuhla, C. Martin, P. J. Moate, P. Niu, N. Peiren, C. Reynolds, S. R. O. Williams, and P. Lund. 2024. Predicting CO₂ production of lactating dairy cows from animal, dietary, and production traits using an international dataset. *J. Dairy Sci.* 107:6771–6784. <https://doi.org/10.3168/jds.2023-24414>.
- López-Paredes, J., I. Gouri, R. Atxaerandio, A. García-Rodríguez, E. Ugarte, J. A. Jiménez-Montero, R. Alenda, and O. González-Recio. 2020. Mitigation of greenhouse gases in dairy cattle via genetic selection: 1. Genetic parameters of direct methane using noninvasive methods and proxies of methane. *J. Dairy Sci.* 103:7199–7209. <https://doi.org/10.3168/jds.2019-17597>.
- Løvendahl, P., V. Milkevych, R. Nielsen, M. Bjerring, C. Manzanilla-Pech, K. Johansen, G. F. Difford, and T. Villumsen. 2024. A data-driven approach to the processing of sniffer-based gas emissions data from dairy cattle. *Comput. Electron. Agric.* 227:1–16. <https://doi.org/10.1016/j.compag.2024.109559>.
- Madsen, J., B. S. Bjerg, T. Hvelplund, M. R. Weisbjerg, and P. Lund. 2010. Methane and carbon dioxide ratio in excreted air for quantification of the methane production from ruminants. *Livest. Sci.* 129:223–227. <https://doi.org/10.1016/j.livsci.2010.01.001>.
- Manzanilla-Pech, C. I. V., D. Gordo, G. F. Difford, P. Løvendahl, and J. Lassen. 2020. Multitrait genomic prediction of methane emissions in Danish Holstein cattle. *J. Dairy Sci.* 103:9195–9206. <https://doi.org/10.3168/jds.2019-17857>.
- Manzanilla-Pech, C. I. V., P. Løvendahl, D. Mansan Gordo, G. F. Difford, J. E. Pryce, F. Schenkel, S. Wegmann, F. Miglior, T. C. Chud, P. J. Moate, S. R. O. Williams, C. M. Richardson, P. Stothard, and J. Lassen. 2021. Breeding for reduced methane emission and feed-efficient Holstein cows: An international response. *J. Dairy Sci.* 104:8983–9001. <https://doi.org/10.3168/jds.2020-19889>.
- Manzanilla-Pech, C. I. V., R. B. Stephansen, G. F. Difford, P. Løvendahl, and J. Lassen. 2022b. Selecting for Feed Efficient Cows Will Help to Reduce Methane Gas Emissions. *Front. Genet.* 13:885932. <https://doi.org/10.3389/fgene.2022.885932>.
- Manzanilla-Pech, G. F., G. F. Difford, P. Løvendahl, R. B. Stephansen, and J. Lassen. 2022a. Genetic (co) variation of methane emissions, efficiency, and production traits in Danish Holstein cattle along and across lactations. *J. Dairy Sci.* 105:9799–9809. <https://doi.org/10.3168/jds.2022.2121>.
- Pedersen, S., and H. Blanes-Vidal. Jørgensen, A. Chwalibog, A. Haeussermann, M.J.W. Heetkamp, A.J.A. Aarnink. 2008. Carbon dioxide production in animal houses: a literature review. *Agric. Eng. Int.: CGIR E journal*, Manus BC 08 008, vol. X (2008)
- Reintke, J., K. Brügemann, T. Yin, P. Engel, H. Wagner, A. Wehrend, and S. König. 2020. Assessment of methane emission traits in ewes using a laser methane detector: genetic parameters and impact on lamb weaning performance. *Arch. Tierzucht* 63:113–123. <https://doi.org/10.5194/aab-63-113-2020>.
- Rey, J., R. Atxaerandio, R. Ruiz, E. Ugarte, O. González-Recio, A. García-Rodríguez, and I. Gouri. 2019. Comparison Between Non-Invasive Methane Measurement Techniques in Cattle. *Animals (Basel)* 9:563. <https://doi.org/10.3390/ani9080563>.
- Sypniewski, M., T. Strabel, and M. Pszczola. 2021. Genetic Variability of Methane Production and Concentration Measured in the Breath of Polish Holstein-Friesian Cattle. *Animals (Basel)* 11:3175. <https://doi.org/10.3390/ani11113175>.
- van Breukelen, A. E., M. N. Aldridge, R. F. Veerkamp, L. Koning, L. B. Sebek, and Y. de Haas. 2023. Heritability and genetic correlations between enteric methane production and concentration recorded by GreenFeed and sniffers on dairy cows. *J. Dairy Sci.* 106:4121–4132. <https://doi.org/10.3168/jds.2022-22735>.
- van Breukelen, A. E., R. F. Veerkamp, Y. de Haas, and M. N. Aldridge. 2024. Genetic parameter estimates for methane emission from breath during lactation and potential inaccuracies in reliabilities assuming a repeatability versus random regression model. *J. Dairy Sci.* 107:5853–5868. <https://doi.org/10.3168/jds.2024-24285>.




Figure 1. Genetic correlations among the CH₄c per visit (avg), sum of max peaks per minute (speaks), number of peaks per minute (npeaks), CH₄p calculated by Madsen, Chagunda, Kjeldsen2, Kjeldsen3 and Tier2, carbon dioxide concentration per visit (CO₂c), milk yield (MY), methane intensity (Mel) and body weight (BW).

NOTES

C. I. V. Manzanilla-Pech, <https://orcid.org/0000-0003-1552-212X>

A. E. van Breukelen, <https://orcid.org/0000-0003-1576-4153>

R. F. Veerkamp, <https://orcid.org/0000-0002-5240-6534>

Y. de Haas, <https://orcid.org/0000-0002-4331-4101>

M. van Pelt, <https://orcid.org/0000-0003-1315-1329>

B. Gredler-Grandl <https://orcid.org/0000-0002-5916-8639>

This article does not contain any studies with human or animal subjects and did not require [IACUC/IRB] approval. This work has received funding from the European Union's Horizon Europe research and innovation programme, grant agreement No 01059609 (Re-Livestock project). The authors also acknowledge Integral approach project "From breeding values to bull selection" for contributing to the funding of this study (BO-43.10-002)